

INFORMED CHOICES ABOUT

COLONOSCOPY

Bowel Preparation Instructions

Very important!

Follow these instructions to clean the stool (poop) out of your colon. Carefully following these instructions will give the best results. If the doctor cannot see the inside of your colon well because of poor cleaning, the test is less useful and very important things can be missed. You may have to repeat the test another time. The risks of the test are greater if the colon is not clean.



Your colonoscopy doctor may give you instructions for preparation that are different. Please follow those if they are different than below.




Step 1. One week before your colonoscopy: Do not eat corn, flax/poppy/ sesame seeds, nuts, fish oils, or medicines containing iron including multivitamins containing iron.

Step 2. A few days before your colonoscopy: Go to any pharmacy and buy the following special laxatives. You do not need a prescription, but they may be kept behind the pharmacy counter. Ask the pharmacist. (Your doctor may suggest different laxatives than the ones below. Please follow your doctor's instructions.)



- One box of Bi-PegLyte preparation. The box contains 2 packets of laxative powder that you will later mix with water to make two 1 L PegLyte solutions. The box also contains 3 bisacodyl tablets.

- Your doctor may ask you to use another laxative such as CoLyte, PegLyte, KleanLyte or Pico-Salax as alternatives to the “Bi-PegLyte.” **Use the laxative your doctor recommends.**



- Some people have a problem with nausea when taking laxatives. You may want to pick up an anti-nausea medicine like Graval. The pharmacist can show you the choices for this.

For more information on colonoscopy visit: mycolonoscopy.ca.

Harminder Singh and John R. Walker, 2023, on behalf of the Optimizing Colonoscopy Team, University of Manitoba, Canada. This material is licensed under a Creative Commons Attribution 4.0 International License. You are free to copy and distribute this material in its entirety as long as it is not altered in any way (no derivative works).



COLONOSCOPY

Step 3. The day before your colonoscopy



- Do not eat any solid foods—until the colonoscopy is finished.
- Drink only clear fluids—until two hours before the time of the colonoscopy. These are liquids that you can see through. See the ‘Fluids that are okay’ list below. Coffee or tea are okay with sweetener but no milk or whitener.
- Try to drink one glass of clear fluids each hour that you are awake.
- Mix the entire contents of one package of laxative powder into 1 liter (32 ounces) of lukewarm water until a clear solution forms. Do not use cold water. Repeat the step with the second package. Do not add anything else (e.g. flavoring, juice). You may want to keep this PegLyte solution in the refrigerator to make it taste better.



Fluids that are okay:

- ✓ clear soups
- ✓ apple juice, **white** grape juice, **white** cranberry juice
- ✓ clear or lemon Gatorade/Powerade
- ✓ Ginger ale, Kool-Aid and water
- ✓ coffee or tea (sugar and sweetener are okay)
- ✓ popsicles (not red or purple)
- ✓ lemon or orange Jello

Food and Fluids that are not okay:

- ✗ no bread, grain or rice
- ✗ no soups with chunks of food
- ✗ no meat, fish, poultry or seafood
- ✗ no fruit or vegetables
- ✗ no milk or dairy products
- ✗ no **red**, **blue** or **purple** liquid
- ✗ no alcoholic drinks
- ✗ Dark colored soft drinks (Pepsi and Coke)

Step 4. The day before the colonoscopy – taking the laxatives.



- At around 2 PM: Take the 3 bisacodyl tablets. Do not take any **antacids** within one hour of taking bisacodyl.
- In the evening at around 8 PM, drink 1 liter of PegLyte solution. Drink about 1 cup (250 mL) every 10 minutes.
- If you feel sick, wait 20 to 30 minutes, then continue to drink the prep.
- You may take Graval (25 to 50 mg) or a similar medicine if the nausea continues. Graval may make you drowsy.

For more information on colonoscopy visit: mycolonoscopy.ca.

Harminder Singh and John R. Walker, 2023, on behalf of the Optimizing Colonoscopy Team, University of Manitoba, Canada. This material is licensed under a Creative Commons Attribution 4.0 International License. You are free to copy and distribute this material in its entirety as long as it is not altered in any way (no derivative works).

COLONOSCOPY

- It is important to take all of the laxatives. The laxatives will cause frequent loose bowel movements. The goal is to have liquid bowel movements, clear to yellow in colour. **Remember, when you drive through a snowstorm it is difficult to see the road. Similarly, your doctor can't get a clear view for the colonoscopy unless your colon is very clean.**
- Your bowel movements should be liquid, clear to yellow in color, which allows your doctor to see your bowel more clearly.



Not ok Not ok Not ok Almost Ready!

Step 5. The day of your colonoscopy – 4-5 hours before the colonoscopy



- **Eat no solid foods, drink only clear fluids.**
- Drink about 1 cup (250 mL) of the second liter of PegLyte solution every 10 minutes until you finish drinking the second liter.
- **STOP DRINKING ALL FLUIDS** 2 hours before your colonoscopy.

Step 6. Have someone take you to the test and someone to take you home from the test



- It is important to have someone with you who can stay to hear the results provided after the test and then take you home. Ask them to write down the results and whether another colonoscopy is suggested in future years. You will feel drowsy after the colonoscopy due to the medications used to relax you. You may not remember things you are told about the results. This means it would be dangerous to drive a car immediately after the test.
- Someone should be with you for up to 24 hours afterwards while the drowsiness and forgetfulness wears off. You will not be able to drive a car.

For more information on colonoscopy visit: mycolonoscopy.ca.

Harminder Singh and John R. Walker, 2023, on behalf of the Optimizing Colonoscopy Team, University of Manitoba, Canada. This material is licensed under a Creative Commons Attribution 4.0 International License. You are free to copy and distribute this material in its entirety as long as it is not altered in any way (no derivative works).

COLONOSCOPY



Questions about your colonoscopy: If you have questions about the preparation for the colonoscopy, or if you run into problems, you are welcome to call the office of the doctor who will do the colonoscopy (unless you were given instructions to call a different number). The staff there are used to handling questions. Most of the offices are open on regular business days from 9 AM to 4 PM. Your pharmacist will be very familiar with your regular medications and the different laxatives used for preparing for the colonoscopy and can also provide helpful information.



Cancellation: If you have to cancel the test, please inform the colonoscopy booking office as soon as possible. Please remember that if you cancel at the last minute or miss the appointment without cancelling, that will lead to wasted time when a colonoscopy could have been done for someone else who is waiting. If you miss or cancel your colonoscopy, there may be a several months wait for a rebooked colonoscopy.

Hence it is important to call as soon as possible to cancel and make another appointment.



Results of your colonoscopy: In addition to the results, you receive just after your colonoscopy you may receive a written report. If you have questions about the results contact the doctor who did the colonoscopy or your family doctor.

COLONOSCOPY

Problem Solving



Problem: You may find it hard to drink all of the liquid. Some people don't like the taste and some have difficulty in drinking the amount needed to clean the colon properly.

Suggestion: Some people find that drinking with a straw helps – so you don't taste it as much. You can rinse your mouth out with a clear drink after each glassful (e.g. ginger ale, 7 Up®, apple juice). If you feel really full, it can help to walk around.

- Cool it to your preferred temperature in the fridge. If you start to feel cold, you can take it out of the fridge or drink some clear warm liquid like broth or herbal tea.



Problem: If your colonoscopy is scheduled for the morning time (say 9 AM) you will have to get up at 4 or 5 AM. Some people find this difficult.

Suggestion: Research shows that taking half of the liquid laxative the evening before the test and half the morning of the test is better for cleaning the bowel and results in better colonoscopies. Remember that people often have to get up early before a long trip also. Plan ahead for some activities you can do early in the morning (radio, music, TV, computer) while taking frequent trips to the toilet.



Problem: Some people feel gassy after the test because of the air used to help do the colonoscopy.

Suggestion: Walking may help with this feeling.

Problem: You have a long drive to reach the colonoscopy facility/hospital.

Suggestion: Most people find that they do not have to make frequent visits to the toilet in the two hours before the colonoscopy. If you have to travel for more than one hour, it may be best to stay somewhere closer to the colonoscopy facility starting the evening before the colonoscopy. You should try to finish your laxative in the morning half to one hour before you leave your home/hotel.

